

Secret
NOFORN



USSR—Current Status of the 1977 Grain Crop

Secret
GC AB 77-003
8 July 1977

Warning Notice
Sensitive Intelligence Sources and Methods Involved
(WNINTEL)


NATIONAL SECURITY INFORMATION
Unauthorized Disclosure Subject to Criminal Sanctions

DISSEMINATION CONTROL ABBREVIATIONS

NOFORN-	Not Releasable to Foreign Nationals
NOCONTRACT-	Not Releasable to Contractors or Contractor/Consultants
PROPIN-	Caution—Proprietary Information Involved
NFIBONLY-	NFIB Departments Only
ORCON-	Dissemination and Extraction of Information Controlled by Originator
REL...-	This Information has been Authorized for Release to ...

Classified by 019641
Exempt from General Declassification Schedule
of E.O. 11652, exemption category:
Sec. 5B(1), (2) and (3)
Automatically declassified on:
date impossible to determine

SECRET
NOFORN



Environment and Resource Analysis Brief

USSR—Current Status of the 1977 Grain Crop

Summary

As of 5 July, we estimate total Soviet grain production at 225 million tons—the same as the current US Department of Agriculture (USDA) projection. This is slightly above last year's record harvest. Wheat production is expected to reach 107 million tons, second only to the 1973 output. USDA puts the wheat harvest at 110 million tons.

Excessive moisture in European USSR has caused harvesting difficulties in several parts of the winter grain area, jeopardizing the bread-making quality of the winter wheat crop.

With 95 percent of the total grain crop yet unharvested and with about two-thirds of the spring grain prior to early-ripening, final outturn is far from certain. Prevailing weather and crop conditions from now through late harvesting in September will have a major impact on the size and quality of the crop.

Note: This paper was produced by the Office of Geographic and Cartographic Research and coordinated with the Office of Economic Research. Comments and questions may be directed to [REDACTED] Code 143, Extension 3748. Date of information 5 July 1977.

25X1A

1
SECRET

SECRET

Total Grain Production

Prospects continue to be good for a second consecutive bumper Soviet grain harvest. We currently project total production at 225 million tons, slightly above last year's record output. (See Chart) We expect wheat to reach about 107 million tons, second only to the 110 million tons produced in 1973; barley output will also be well above average. Only corn will fall below the average production of recent years.


In the midst of an excellent winter grain harvest, the key to achieving a new record requires maintaining adequate soil moisture levels in the drought-prone spring grain area, especially in the spring wheat belt of Kazakhstan and West Siberia. (See Map 1) The short range weather forecast indicates a stationary weather system that will provide adequate rainfall in most of the spring grain area through mid-July.

Weather conditions from now till September will largely determine the size and quality of this year's grain harvest. As of 1 July, less than 5 percent of the total grain crop had been harvested and some two-thirds of the spring grains had not ripened.

Status of the Winter Grains


The 1977 winter grain crop will be about 8½ million tons greater than the previous record and underpins prospects for a bumper total grain harvest. As of early July, we estimate output of winter grains at 72 million tons—some 2 million tons higher than our initial projection in late May.¹ The upward revision reflects slightly better yields from both the winter rye and winter barley, crops that account for about ⅓ of the harvested winter grain acreage. This year's production marks a sharp improvement over output of winter grains in 1976. Of the total winter grain harvest, we expect winter wheat to account for at least 57 million tons, compared with 45 million tons last year. (See Table 1)

¹ See USSR—*Current Status of 1977 Grain Crops*, 27 May 1977, GC AB 77-002. S.



SECRET

USSR: Major Spring and Winter Grain Area



573079 5-77 (542601)

■ Spring and winter grains

□ Spring grains

----- Economic region boundary

SECRET

SECRET

Table 1
USSR: Grain Production ¹

	Million tons							
	1970	1971	1972	1973	1974	1975	1976	Estimate 1977
TOTAL	186.8	181.2	168.2	222.5	195.6	140.1	223.8	225
Winter Grains ²	57.5	63.0	40.6	63.5	62.5	48.6	60.9	72
Wheat	42.2	47.8	29.4	49.4	44.7	36.6	44.6	57
Rye	13.0	12.8	9.6	10.8	15.2	9.0	14.0	13
Spring Grains ³	129.3	118.2	127.6	159.0	133.1	91.2	162.9	153
Wheat	57.4	51.1	56.6	60.5	39.2	29.5	52.3	50
Barley	35.8	32.3	35.1	51.7	51.6	32.8	67.1	60

¹ Because of rounding, components may not add to totals shown.

² In addition to winter wheat and rye, winter grains include winter barley.

³ In addition to spring wheat and barley, spring grains include oats, corn for grain, pulses, millet, rice, and buckwheat.

In the fall of 1976, winter grains—wheat, rye, and barley—were sown on 38½ million hectares, the largest sown area since 1968 and about 1 million hectares greater than the previous fall.

Most of the winter grains went into dormancy in very good condition; soil moisture reserves were uniformly excellent. Published Soviet accounts indicate that winterkill was light except in parts of the Non-Chernozem Zone and North Caucasus. Areas sown to rye and barley suffered the most damage. Imagery revealed only slight damage to winter wheat in the Ukraine. We expect that 83 percent of the winter grain acreage will be harvested this year, compared with the ten-year average of less than 80 percent.

Harvesting, which began in late June in the southern region around the Black Sea, is especially difficult this year because of heavy June rainfall. (See Map 2) The U.S. Department of Agriculture (USDA) winter grain team reported extensive lodging in the eastern Ukraine. A Soviet official confirms that 80% of the combines were adjusted for harvesting flattened crops.²


Wet conditions are also delaying mechanized field operations, such as haymaking, and in turn, these operations are overlapping the early wheat harvest in parts of the North Caucasus. Heavy weed growth—another consequence of the excessive rainfall—will add to harvesting difficulties.

Unless a drying trend develops by mid-July, the milling and bread-making quality of winter wheat will be substantially reduced by excess moisture and heavy mildew. Although Soviet grain drying capacity has expanded in recent years, facilities will face a stiff test in coping with this year's crop. Storing grain with a moisture

² Lodging describes a condition resulting when, because of rain and wind, grain stalks bend or break and form a flattened or tangled mass that is difficult to harvest. It generally occurs during the later stages of crop development—when the grains are tallest and weighted down with mature heads—and when plant growth has been especially lush. The extent of the losses associated with lodging depends partly on the stage of the plants' development when lodging occurs, the type of harvesting method employed, and weather conditions during harvesting. Production losses due to post-maturity lodging are minimized in the USSR by the use of hand harvesting.

SECRET

USSR: Area of Excess Moisture as of 1 July 1977



SECRET

SECRET

Table 4
USSR: Harvested Area

	Million hectares				
	1973	1974	1975	1976	Estimate 1977
Winter Grain	26.9	29.9	29.2	27.4	32
Spring Grain	99.8	97.3	98.7	100.3	97 ¹
TOTAL	126.7	127.2	127.9	127.7	129

¹ Reflects normal abandonment of about 2 million hectares.

Early spring soil moisture was above normal in most of European USSR with the exception of the Southern Urals and parts of the Volga valley. Although both of the latter areas were dry during optimum seeding in late May, subsequent rainfall significantly improved both soil moisture levels and plant vigor.


During June and early July unusually heavy rains occurred throughout European USSR from the Baltics to the North Caucasus. Apart from some localized flooding in the Ukraine, we expect the overall impact of above normal precipitation on this years crop to be unusually favorable. Yields on both winter and spring grains in the Southern Ukraine adjacent to the Black Sea could be as much as 10 percent above their historical maximum.

In the spring, soil moisture in northern Kazakhstan and West Siberia was above normal and substantially better than last year. Parts of northern Kazakhstan experienced a reduction in surface soil moisture in early May, but improved markedly as a result of widespread local showers during June. In West Siberia, favorable crop conditions have existed since early spring and show no signs of deteriorating. Infra-red (IR) imagery taken in mid-June depicts excellent plant vigor levels, indicating better than average yield potential. (See Imagery)

SECRET


SECRET

Landsat Imagery, Mid-June 1976



Bright infra-red reflectance (IR) in the spring wheat region of West Siberia depicts equally good crop vigor levels for the last two years.

Landsat Imagery, Mid-June 1977



573425 7-77 CIA

SECRET
NOFORN

* Note: Due to methods used in processing red return may be slightly different.

Area of imagery

SECRET

Secret

Secret